

**VLA-5 is expressed by mouse and human long-term repopulating hematopoietic cells and mediates adhesion to extracellular matrix protein fibronectin.**

Johannes C.M. van der Loo, ... , Ikunoshin Kato, David A. Williams

*J Clin Invest.* 2001;107(10):1203-1203. <https://doi.org/10.1172/JCI3687C1>.

**Corrigendum**

*J. Clin. Invest.* 102:1051–1061 (1998) The peptide concentration on the x-axis of figure 2 should read pmol/cm<sup>2</sup>. The concentrations mentioned in the text and legends, which were based on this figure, should also read pmol/2. The corrected figure and legend appear below. Figure 2 Plastic adherence of FN peptides. Effect of the coating concentration on the absorption of recombinant FN fragments to nontissue culture-treated plastic (A) and on the subsequent binding of mouse BM cells or human HEL cells (B). Bound FN was detected by ELISA (average of 4 wells/concentration) using antibody FN12-8 (A), which is specific for the CELL-binding domain of FN. (A) Fragment H-271 (filled squares), C-274 (open squares), CH-271 (filled circles), H-296 (open circles), and CH-296 (filled triangles). For clarity, error bars have been omitted. (B) Binding of human HEL cells (squares) or low-density mouse BM cells (circles) to plates coated with H-271 (open symbols) or CH-296 (closed symbols).

**Find the latest version:**

<https://jci.me/3687C1/pdf>



**Dynamin is involved in human epithelial cell vacuolation caused by the *Helicobacter pylori*-produced cytotoxin VacA**

Junko Suzuki, Hirohide Ohnishi, Hiroshi Shibata, Akihiro Wada, Toshiya Hirayama, Taroh Iiri, Namiki Ueda, Chiho Kanamaru, Tomohiro Tsuchida, Hirosato Mashima, Hiroshi Yasuda, and Toshiro Fujita

*J. Clin. Invest.* **107**:363-370 (2001).


During the preparation of this manuscript for publication, an error was introduced into the list of authors. The corrected information appears above.

**VLA-5 is expressed by mouse and human long-term repopulating hematopoietic cells and mediates adhesion to extracellular matrix protein fibronectin.**

Johannes C.M. van der Loo, Xiangli Xiao, Doug McMilin, Kimikazu Hashino, Ikunoshin Kato, and David A. Williams

*J. Clin. Invest.* **102**:1051-1061 (1998).

The peptide concentration on the x-axis of figure 2 should read pmol/cm<sup>2</sup>. The concentrations mentioned in the text and legends, which were based on this figure, should also read pmol/cm<sup>2</sup>. The corrected figure and legend appear below.



**Figure 2**

Plastic adherence of FN peptides. Effect of the coating concentration on the absorption of recombinant FN fragments to nontissue culture-treated plastic (**A**) and on the subsequent binding of mouse BM cells or human HEL cells (**B**). Bound FN was detected by ELISA (average of 4 wells/concentration) using antibody FN12-8 (**A**), which is specific for the CELL-binding domain of FN. (**A**) Fragment H-271 (filled squares), C-274 (open squares), CH-271 (filled circles), H-296 (open circles), and CH-296 (filled triangles). For clarity, error bars have been omitted. (**B**) Binding of human HEL cells (squares) or low-density mouse BM cells (circles) to plates coated with H-271 (open symbols) or CH-296 (closed symbols).